



Set up (M^4, g)
 $\Sigma^2 \subseteq M^4$ $M - \Sigma$ smooth

Coordinates

local (y_1, y_2, x_1, x_2)

$y_1 = y_2 = 0$ on Σ

polar (p, θ, x_1, x_2)



Edge - Cone Metric

Normal bundle near Σ : $N\Sigma \cong \Sigma^2 \times \mathbb{R}^2$
 $dp^2 + \beta^2 p^2 d\theta^2 + g_\Sigma(x)$

S' -connection $S' \hookrightarrow N\Sigma \xrightarrow{\pi} \Sigma$

Model cone metric

$$\begin{aligned} \bar{g} &= dp^2 + \beta^2 p^2 (d\theta + v)^2 + g_\Sigma \\ &= dp^2 + \beta^2 p^2 (d\theta + \underbrace{u_j(x) dx^j}_{\text{smooth conn. 1-form}})^2 + w_{jk} dx^i dx^k \end{aligned}$$

Edge-cone metric

$$g = \bar{g} + \underbrace{\rho^{1+\epsilon} h}_{\text{perturbation}} \quad \text{on } M \setminus \Sigma$$

perturbation, h symmetric 2-tensor

Einstein edge-cone metric: g Einstein on $M \setminus \Sigma$

e.g. $\beta < \frac{1}{3}$ Kähler-Einstein

$\beta = \frac{1}{p}$, $p \geq 2$, quotient of non-singular Einstein $X/\mathbb{Z}_p$ w/ codim 2 fixed points

4D Hitchin-Thorpe inequality

• smooth compact Einstein

$$2\chi(M) \pm 3\tau(M) \geq 0$$

pf. Gauss-Bonnet type integrals

$$\chi(M) = \frac{1}{8\pi^2} \int_M \left(\frac{\sigma^2}{24} + |W|^2 - \frac{|\text{Ric}^\circ|^2}{2} \right) d\mu$$

$$\tau(M) = \frac{1}{12\pi^2} \int_M (|W_+|^2 - |W_-|^2) d\mu$$

$$\Rightarrow \text{LHS} = \frac{1}{4\pi^2} \int_M \left(\frac{\sigma^2}{24} + 2|W_\pm|^2 - \frac{|\text{Ric}^\circ|^2}{2} \right) d\mu \quad \text{Einstein}$$

Correction needed for χ and τ w/ edge-cone singularity

- edge-cone singularity

$$2\chi(M) \pm 3\tau(M) \geq (1-\beta) \left(2\chi(\Sigma) \pm (1+\beta)[\Sigma]^2 \right)$$

where M^4 smooth compact

Σ^2 smooth embedded compact oriented

(M, Σ) admit Einstein edge cone metric g
w/ angle $2\pi\beta$

Thm 2.1 & 2.2

$$\chi(M) - (1-\beta)\chi(\Sigma) = \frac{1}{8\pi^2} \int_M \left(\frac{s^2}{24} + |W|^2 - \frac{|\mathring{Ric}|^2}{2} \right) d\mu$$

$$\tau(M) - \frac{1}{3}(1-\beta^2)[\Sigma]^2 = \frac{1}{12\pi^2} \int_M (|W_+|^2 - |W_-|^2) d\mu$$

↑ self-intersection of
cohomology class in $H^2(M) \stackrel{PD}{\cong} H_2(M)$

$$= \langle [\Sigma] \cup [\Sigma], [M] \rangle$$

$$2(1-\beta)\chi(\Sigma) \pm (1-\beta^2)[\Sigma]^2$$

$$\Rightarrow 2\chi(M) \pm 3\tau(M) - \boxed{\text{def}_{\pm}(\Sigma, \beta)}$$

$$= \frac{1}{4\pi^2} \int_M \left(\frac{s^2}{24} + 2|W_{\pm}|^2 - \frac{|\mathring{Ric}|^2}{2} \right) d\mu$$

Step 1. Metric-independent

Lemma 2.1 it suffices to check (*) for one edge-cone metric

pf. Ref of edge cone metric controls decay rate of the integrand.

Let $\dot{g} = g' - g$ w/ same angle

$$g_t = g + t\dot{g} + O(t^2)$$

edge-cone $g = \bar{g} + \rho^{1+\varepsilon}h$ almost smooth so ∂g smooth + $O(\rho^\varepsilon)$, $\partial^2 g \sim O(\rho^{-1+\varepsilon})$

$$\Rightarrow |R| \sim O(\rho^{-1+\varepsilon})$$

$$\Rightarrow |\Phi_t|, |\Psi_t| \leq |R| \cdot |\nabla \dot{g}| \quad C\rho^{-1+\varepsilon}$$

$$\left| \frac{d}{dt} \int_{M_\delta} \Phi_t \right| = \int_{\partial M_\delta} \Phi_t \leq C_1 \delta^{-1+\varepsilon}$$

$$\int \quad = C_2 \delta^\varepsilon \xrightarrow{\delta \rightarrow 0} 0$$

Gauss-Bonnet integrand

Step 2 Simplification to a Ricci form

- Let $\Upsilon = \left(\frac{s^2}{24} + 2|W_+|^2 - \frac{|Ric|^2}{2} \right) d\mu$
 g_0 smooth metric g edge-cone metric

Then

$$\int_M \Upsilon_g - \boxed{\int_M \Upsilon_{g_0}} = \int_M \boxed{\Upsilon_g - \Upsilon_{g_0}} = -4\pi^2 \text{def}_+(\Sigma, \beta)$$

$= 4\pi^2 (2\chi + 3\tau)(M)$ ↪ support in a tubular neighbourhood V of Σ

- Kähler $\Rightarrow |W_+|^2 = \frac{s^2}{24}$

$$\Upsilon = \frac{1}{2} \left(\frac{s^2}{4} - |Ric|^2 \right) d\mu = \rho \wedge \rho$$

↗ Ricci form

supp of $\rho \wedge \rho$ zero measure sing. set.

$$\text{LHS} = \int_{V-\Sigma} \rho_g \wedge \rho_g - \rho_{g_0} \wedge \rho_{g_0}$$

$$\stackrel{NTS}{=} -4\pi^2 (1-\beta) \left(2\chi(\Sigma) + (1+\beta)[\Sigma]^2 \right)$$

Step 3. Prove (*) for a special case

Prop 2.1 proves (*) by constructing a specific model metric and compute the defect term explicitly.

- construct a rotationally symmetric Kähler metric

$$\begin{array}{ccc} \nearrow & \Phi(\zeta, \bar{\zeta}) = F(|\zeta|^2) = F(t) & \nwarrow \\ \text{Kähler potential} & & \text{auxiliary func.} \end{array}$$

$$\omega = i\partial\bar{\partial} F(|\zeta|^2) = i f(|\zeta|^2) d\zeta \wedge d\bar{\zeta}$$

local Kähler form

$$\text{Def. } \frac{dF}{dt} = \frac{1}{t} \int_0^t f(x) dx, \quad f(x) = \begin{cases} \frac{1}{\beta} & t \leq \frac{1}{2} \\ t^{\beta-1} & t \geq 1 \end{cases}$$

f controls cone angle

ω normal bundle $E \rightarrow \Sigma$

$$\begin{array}{l} \omega = \lambda \omega^* \alpha + i\partial\bar{\partial}(\beta^{-2} t^\beta) \\ \omega_0 = \lambda \omega^* \alpha + i\partial\bar{\partial} F(t) \end{array} \rightsquigarrow \left. \begin{array}{l} g \\ g_0 \end{array} \right\} \text{agree for } t \geq 1$$

$\underbrace{\hspace{1.5cm}}_{\text{form on } E} \quad \text{area form of } \Sigma$

locally

$$\omega = it^{\beta-1} d\zeta \wedge d\bar{\zeta} + i(\lambda - \frac{\kappa}{\beta} t^\beta) dz \wedge d\bar{z} + O(|z|)$$

$$\omega_0 = if(t) d\zeta \wedge d\bar{\zeta} + i(\lambda - \kappa t F'(t)) dz \wedge d\bar{z} + O(|z|)$$

$\uparrow$
normal fiber $E \ni (x, \zeta)$

$$\int_{V-\Sigma} \rho g^2 - \rho g_0^2 \stackrel{\text{Stoke}}{=} - \int_{S_\varepsilon} \phi \wedge (2\rho_0 + d\phi)$$

where $\rho - \rho_0 = d\phi$ / $\phi = -\delta\theta \log\left(\frac{V}{V_0}\right)$

can compute explicitly

ε width of tubular neighbourhood

$$= 4\pi^2(1-\beta)(2\chi(\Sigma) + (1+\beta)[\Sigma]^2) + \mathcal{O}(\varepsilon^{2\min(1,\beta)})$$

then let $\varepsilon \rightarrow 0$

Remark. If Σ non-oriented, but M oriented, then we need to pass the computation of $\chi(\Sigma)$ to the oriented bouble cover.

Prop 2.2 (M, J) almost complex $2n$ -manifold
orientation induced by J
 Σ totally real w.r.t J

$$\tau(M) + \frac{1}{3}(1-\beta^2)\chi(\Sigma) = \frac{1}{12\pi^2} \int_M (|W_+|^2 - |W_-|^2) d\mu$$

§ 3. G -index proof

$$M = X/G$$

Set up

finite G acting on compact oriented X .

preserving the metric & orientation

Let $x \in X^g$, fixed point set of g , the linearized action decomposes as rotations.

$$g|_{T_x X} = \begin{pmatrix} R(\alpha) & 0 \\ 0 & R(\beta) \end{pmatrix}$$

$$H^2(X; \mathbb{R}) = H^+ \oplus H^-$$

$\forall g \in G$, g_* induced action on $H^2(M; \mathbb{R})$

$$\text{Set } \tau(g, X) = \text{tr}(g_*|_{H^+}) - \text{tr}(g_*|_{H^-})$$

Orbifold signature

$$g=1 \quad \tau(1, X) = \text{signature}$$

$$g \neq 1 \quad X^g = \text{fixed point set of } g$$

$$= \bigsqcup_{\text{isolated}} x_j \cup \bigsqcup \hat{\Sigma}_k$$

$\alpha, \beta \neq 0$

compact surfaces
one of α or $\beta = 0$

$$\Rightarrow \tau(g, X) = - \sum_j \cot \frac{\alpha_j}{2} \cot \frac{\beta_j}{2} + \sum_k \left(\csc^2 \frac{\theta_k}{2} \right) [\Sigma_k]^2$$

Averaging over G yields the orbifold signature

$$\tau(M) = \frac{1}{|G|} \left(\tau(X) + \sum_{g \neq 1} \tau(g, X) \right)$$

$\beta = \frac{1}{p} \quad p \in \mathbb{Z}_{\geq 0}$

Apply this to $G = \mathbb{R}_p$ and use $\sum \csc^2(ka/p) = \frac{p^2-1}{3}$

$$\Rightarrow \tau(M) - \frac{1}{3} \left(1 - \frac{1}{p^2}\right) [\Sigma]^2 = \frac{1}{p} \tau(X)$$

$$= \frac{1}{12\pi^2} \int_M |W_+|^2 - |W_-|^2 d\mu$$

§4. Einstein

Since $2\chi(M) \pm 3\tau(M) - \text{def}_{\pm}(\Sigma, \beta)$

$$= \frac{1}{4\pi^2} \int_M \left(\frac{s^2}{24} + 2|W_{\pm}|^2 - \frac{|\text{Ric}|^2}{2} \right) d\mu$$

we know Einstein $\Rightarrow$ RHS ≥ 0 so

$$2\chi(M) \pm 3\tau(M) \geq \text{def}_{\pm}(\Sigma, \beta) \quad (\star)$$

w/ equality $\Leftrightarrow s=0$ and $W_{\pm} \equiv 0$

$\Leftrightarrow \Lambda^{\pm}$ flat

(Einstein) $\Leftrightarrow$ locally hyper Kähler

(4D) $\Leftrightarrow$ Ricci flat + locally Kähler

• small angle: $\beta_i \rightarrow 0$ Coro 4.2

$$2\chi(M) \pm 3\tau(M) \geq 2\chi(\Sigma) \pm [\Sigma]^2$$

• large angle: obstruction to Einstein metric

$$\frac{1}{\beta^2} (\star^{\pm}) \quad \text{let } \beta \rightarrow \infty \Rightarrow \begin{cases} 0 \geq -[\Sigma]^2 \\ 0 \geq [\Sigma]^2 \end{cases} \Rightarrow \boxed{[\Sigma]^2 = 0}$$

$$\frac{1}{4\beta} ((\star^+) + (\star^-)) \Rightarrow \boxed{0 \geq -\chi(\Sigma)}$$

§ 5.

- families of gravitational instanton
self-dual: $W_- = 0$
complete, non-compact, Ricci-flat M^4 .
- hyperbolic ansatz
 $U \subseteq \mathbb{H}^3$ 3D hyperbolic space

$V: U \rightarrow \mathbb{R}$ harmonic w.r.t. hyperbolic metric
defines a connection Θ 1-form of a principal
 $U(1)$ -bundle via $d\Theta = *dV$.

Set $g_0 = Vh + V^{-1}\Theta^2$ on $\begin{array}{c} P \\ \downarrow \\ U \end{array}$

- Consider potential $V = \beta^{-1} + \sum_{j=1}^n G_{p_j}$
 $\beta > 0$ const.
 superposition of Green's functions in $\mathbb{H}^3$
 p_j distinct pts.

By conformal rescaling $g = \beta (\text{sech}^2 x) g_0$
 $\uparrow$
 dist from arbitrary base pt in $\mathbb{H}^3$.

• compactification

$$M = \mathcal{P} \cup \Sigma \cup \{\hat{p}_j\} \cong n \mathbb{CP}^2 = \mathbb{CP}^2 \# \dots \# \mathbb{CP}^2$$

interior

added 2-sphere comes from sphere at infinity
"cone singularity"

added to compactify
fibre over p_j "smooth"

Near p_j . $V \sim e^{-2\alpha}$,

$g_0 \sim$ Gibbons-Hawking near isolated center
 $\hat{p}_j$ corresponds to non singular nut

Hitchin-Thorpe says

$$\frac{1}{12\pi^2} \int_M |W_+|^2 d\mu = \tau(M) - \frac{1}{3} (1 - \beta^4) [\Sigma]^2 = \frac{n(2 + \beta^2)}{3}$$

Example: $n=1$ $V = \beta^{-1} + (e^{2\alpha} - 1)^{-1}$

$$\tilde{g} = \frac{1}{4\beta} \left((2 - \beta) \cosh \alpha + \beta \sinh \alpha \right)^{-2} g_0$$

is Einstein w/ const = $\frac{3}{2} \beta^2 (2 - \beta)$ $0 < \beta < 2$

$\beta = 1$ Fubini-Study on $\mathbb{CP}^2$

$\beta \rightarrow 0$ Taub-NUT on $\mathbb{R}^4$

$\beta \rightarrow 2$ Eguchi-Hanson on $T\mathbb{S}^2$.