

Last time Calc I Review

This time Integration by parts

Motivation

using Calc I knowledge (FTC) we know the
antiderivative of x is $\frac{1}{2}x^2 + C$
 $\cos x$ $\sin x$

It is natural to ask what is the antiderivative
of $\ln x$?

$\tan x$?

To solve this problem we need a new tool.

Recall product rule

$$(uv)' = u'v + uv'$$

if we integrate on both sides w.r.t. x .

$$\begin{aligned}\Rightarrow \int uv \, dx &= \int uv' \, dx + \int u'v \, dx \\ &= \int u \, dv + \int v \, du\end{aligned}$$

rearrange

\Rightarrow


$$\int u \, dv = uv - \int v \, du$$

Example 1 compute $\int \ln x \, dx$

take $u = \ln x$ $v = x$

$$\begin{aligned}\int \ln x \, dx &= x \ln x - \int x \, d \ln x \\ &= x \ln x - \int x \frac{1}{x} \, dx \\ &= x \ln x - x + C\end{aligned}$$

Intuition



NB. There's a list of choices called LIATE rule tells which function to choose as u first
However in general there's no easy way to tell immediately which function to take as u or v .

L \ln log
I \arcsin \arccos \arctan ...
A algebraic x $1+x^2$
T \sin \cos ...
E e^x

Example 2 $\int \underbrace{\arctan x}_u \underbrace{dx}_v$

$$= x \arctan x - \int x d \arctan x$$

$$= x \arctan x - \int \frac{x}{1+x^2} dx$$

use substitution law

$$= \frac{1}{2} \int \frac{d(x^2+1)}{x^2+1} = \frac{1}{2} \ln(1+x^2) + \tilde{C}$$

$$= x \arctan x - \frac{1}{2} \ln(1+x^2) + C$$

Example 3 $\int x e^x dx = \int \underbrace{x}_u \underbrace{de^x}_v$

$$= x e^x - \int e^x dx$$

$$= x e^x - e^x + C$$

Step

1. Choose function u .
2. The choice of v depends on u .

That is, in order to match the integration by parts formula, say we are computing $\int f(x) dx$

Then we claim $dv = \frac{f'(x)}{u(x)} dx$, so that

$$\int f dx = \int u \cdot \frac{f}{u} dx = \int u dv$$

$$\begin{aligned}
 \text{Example 4} \quad \int \frac{x^3}{\sqrt{1+x^2}} dx &= \int x^2 \underbrace{\frac{x}{\sqrt{1+x^2}} dx}_{\frac{1}{2} d(x^2+1)} \\
 &= \frac{\frac{1}{2} d(x^2+1)}{\sqrt{1+x^2}} = d\sqrt{1+x^2} \\
 &= \int \underbrace{x^2}_u \underbrace{d\sqrt{1+x^2}}_v \\
 &= x^2 \sqrt{1+x^2} - \int \sqrt{1+x^2} dx^2 = d(x^2+1) \\
 &= x^2 \sqrt{1+x^2} - \frac{2}{3} (1+x^2)^{\frac{3}{2}} + C
 \end{aligned}$$

NB. There's more than one way to solve Ex 4.

$$\text{Example 4'} \quad \int \frac{x^3}{\sqrt{1+x^2}} dx$$

substitution rule $u = 1+x^2 \quad du = 2x dx$
 $x^3 dx = x^2 \cdot (x dx) = \frac{1}{2} (u-1) du$

$$\begin{aligned}
 \int \frac{x^3}{\sqrt{1+x^2}} dx &= \int \frac{\frac{1}{2} (u-1) du}{\sqrt{u}} \\
 &= \frac{1}{2} \int \underbrace{\frac{u}{\sqrt{u}} - \frac{1}{\sqrt{u}}}_{= u^{\frac{1}{2}} - u^{-\frac{1}{2}}} du \\
 &= \frac{1}{3} u^{\frac{3}{2}} - u^{\frac{1}{2}} + C \\
 &= \frac{1}{3} (1+x^2)^{\frac{3}{2}} - (1+x^2)^{\frac{1}{2}} + C
 \end{aligned}$$

Last time Integration by parts

$$\int u dv = uv - \int v du$$

This time: we focus on a particular type of integral

$$\int \sin^m x \cos^n x dx \quad \text{⊗}$$

↑ integers

Tools to solve ⊗

(A) substitution rule

(B) trig identity

(C) double angle formulae

$$\cos^2 x + \sin^2 x = 1$$

$$\sin(2x) = 2 \sin x \cdot \cos x$$

$$\cos(2x) = \cos^2 x - \sin^2 x = 2 \cos^2 x - 1$$

$$= 1 - 2 \sin^2 x$$

! Remember these equations

Note that (D) implies

$$\sin^2 x = \frac{1 - \cos 2x}{2}$$

$$\cos^2 x = \frac{1 + \cos 2x}{2}$$

We'll start by looking at the following examples.

$$\begin{aligned}
\text{Example 1. } & \int \sin^3 x \cos^2 x \, dx \\
&= \int \underbrace{\sin^2 x}_{1 - \cos^2 x} \cos^2 x \underbrace{\sin x \, dx}_{\text{substitution } u = \cos x} \\
&= \int (1 - u^2) u^2 (-du) \quad du = -\sin x \, dx \\
&= \int u^2 - u^4 \, du \\
&= \frac{u^3}{3} - \frac{u^5}{5} + C \\
&= \frac{\cos^3 x}{3} - \frac{\cos^5 x}{5} + C
\end{aligned}$$

In general, if m or n is odd we can use substitution rule $u = \cos x, \sin x$.

$$\begin{aligned}
\text{Example 2 } & \int \cos^2 x \sin^2 x \, dx \\
&= \int \frac{1 + \cos 2x}{2} \cdot \frac{1 - \cos 2x}{2} \, dx \\
&= \frac{1}{4} \int 1 - \cos^2 2x \, dx \\
&= \frac{1}{4} \int 1 - \frac{1 + \cos 4x}{2} \, dx \\
&= \frac{1}{8} \int 1 - \cos 4x \, dx \\
&= \frac{1}{8} x - \frac{1}{32} \sin 4x + C
\end{aligned}$$

If m and n both are even we can use trig identity and double angle formulae.

Similarly we can compute

$$\int \tan^m x \sec^n x \, dx$$

Tools to solve \otimes

(A) substitution rule

(B) trig identity $\sec^2 x = 1 + \tan^2 x$!

(C) double angle formulae

Recall

$$(\tan x)' = \sec^2 x \quad (\sec x)' = \sec x \tan x$$

Example 3. $\int \tan x \sec^4 x \, dx$

$$= \int \tan x \sec^2 x \sec^2 x \, dx$$

$$= \int \tan x (1 + \tan^2 x) \sec^2 x \, dx$$

$$= \int u(1 + u^2) \, du \quad \leftarrow u = \tan x$$

$$= \frac{1}{2} u^2 + \frac{1}{4} u^4 + C$$

$$= \frac{1}{2} \tan^2 x + \frac{1}{4} \tan^4 x + C$$

Another way to compute this

$$\text{Take } u = \sec^2 x$$

$$du = 2 \sec^2 x \cdot \tan x \, dx$$

$$\text{Then } \int \tan x \sec^4 x \, dx$$

$$= \int u \cdot \frac{1}{2} du$$

$$= \frac{u^2}{4} + \tilde{C}$$

$$= \frac{\sec^4 x}{4} + \tilde{C}$$

$$\begin{aligned} \textcircled{*} \left\{ \begin{aligned} &= \frac{1}{4} (1 + \tan^2 x)^2 + \tilde{C} \\ &= \frac{\tan^4 x}{4} + \frac{\tan^2 x}{2} + \underbrace{\frac{1}{4} + \tilde{C}}_{= C} \end{aligned} \right. \end{aligned}$$

⊗ Here I'm checking these two methods give the same solution. You don't need to write these in homework.

In this section we consider the contains square roots of the form

$$\sqrt{a^2 - x^2} \quad \sqrt{x^2 + a^2} \quad \sqrt{x^2 - a^2}$$

We will make trig substitutions

$$\sqrt{a^2 - x^2} \quad \sqrt{x^2 + a^2} \quad \sqrt{x^2 - a^2}$$

$$x = a \sin \theta \quad a \tan \theta \quad a \sec \theta$$

$$\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \quad \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \quad \left[0, \frac{\pi}{2}\right) \text{ or } \left[\pi, \frac{3\pi}{2}\right)$$

$$\cos x \geq 0 \quad \sec x \geq 0 \quad \tan x \geq 0$$

Note that we can use trig identities to remove square roots.

$$\begin{aligned} \text{e.g. } \sqrt{a^2 - x^2} & \stackrel{x = a \sin \theta}{=} \sqrt{a^2 - a^2 \sin^2 \theta} \\ & = \sqrt{a^2 \cos^2 \theta} \\ & = |a \cos \theta| \end{aligned}$$

NB. We have to specify range of θ so that we can remove $| |$.

Example 1. $\int \sqrt{9-x^2} dx$

Take $x=3\sin\theta$ $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$

$$= \int \sqrt{9-(3\sin\theta)^2} d(3\sin\theta)$$

$$= \int \underbrace{|3\cos\theta|}_{=3\cos\theta} \cdot 3\cos\theta d\theta$$

Note that $\cos\theta > 0$ for $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$

$$= 9 \int \cos^2\theta d\theta$$

$$= 9 \int \frac{1+\cos 2\theta}{2} d\theta$$

$$= \frac{9}{2} \int 1+\cos 2\theta d\theta$$

$$= \frac{9}{2} \theta + \frac{9}{4} \sin 2\theta + C$$

$$2\sin\theta\cos\theta$$

$$= \frac{9}{2} (\theta + \sin\theta\cos\theta) + C$$

$$= \frac{9}{2} \left(\arcsin \frac{x}{3} + \frac{x}{3} \cdot \sqrt{1-\frac{x^2}{3^2}} \right) + C$$

$$= \frac{9}{2} \arcsin \frac{x}{3} + \frac{x\sqrt{9-x^2}}{2} + C$$

Example 2 $\int \frac{1}{x^2 \sqrt{x^2+4}} dx$

Take $x = 2 \tan \theta \quad -\frac{\pi}{2} < \theta < \frac{\pi}{2}$

$$= \int \frac{d(2 \tan \theta)}{4 \tan^2 \theta \cdot \sqrt{4 \tan^2 \theta + 4}}$$

$$= \int \frac{2 \sec^2 \theta d\theta}{4 \tan^2 \theta \sqrt{4 \sec^2 \theta}}$$

$$= \frac{1}{4} \int \frac{\sec^2 \theta d\theta}{\tan^2 \theta \cdot \sec \theta}$$

$$= \frac{1}{4} \int \frac{\sec \theta}{\tan^2 \theta} d\theta$$

$$\frac{1}{\cos \theta} \cdot \frac{\cos^2 \theta}{\sin^2 \theta} = \frac{\cos \theta}{\sin^2 \theta}$$

$$= \frac{1}{4} \int \frac{\cos \theta}{\sin^2 \theta} d\theta$$


$$\frac{d(\sin \theta)}{\sin^2 \theta}$$

$$= -\frac{1}{4} \frac{1}{\sin \theta} + C$$

$$= -\frac{\sqrt{4+x^2}}{4x} + C$$

Recall $\tan \theta = \frac{x}{2}$

$$\Rightarrow \sin \theta = \frac{x}{\sqrt{4+x^2}}$$



$$\text{Example 3. } \int \frac{x}{\sqrt{3-2x-x^2}} dx$$

$$= -(x^2+2x+1-1)+3$$

$$= 4-(x+1)^2$$

Take $u = x+1$

$$= \int \frac{u-1}{\sqrt{4-u^2}} d(u-1)$$

Take $u = 2\sin\theta$

$$= \int \frac{2\sin\theta - 1}{\sqrt{4-4\sin^2\theta}} d(2\sin\theta) \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

$$= \int \frac{2\sin\theta - 1}{|2\cos\theta|} (2\cos\theta) d\theta$$

$$= \int 2\sin\theta - 1 d\theta$$

$$= -2\cos\theta - \theta + C$$

$$= -\sqrt{4-u^2} - \arcsin \frac{u}{2} + C$$

$$= -\sqrt{3-2x-x^2} - \arcsin\left(\frac{x+1}{2}\right) + C$$

Last time integral containing $\sqrt{\quad}$
This time integrating rational functions

Defining rational functions

$$R(x) = \frac{P(x)}{Q(x)} \quad \text{where } P, Q \text{ are polynomials}$$

$$\text{e.g. } \frac{1}{x+1}, \quad \frac{2x+1}{x^2+4x+3}, \quad \frac{x^4-2}{(x-1)(x^2+1)}$$

If $\deg P < \deg Q$, we call R a proper rational function

$$\text{e.g. } \frac{1}{x^4-1} \quad \frac{x}{x^2+2} \quad \text{proper}$$

$$\frac{x^2}{x^2+2} \quad \frac{x^4}{(x-1)^2} \quad \text{improper}$$

How to solve $\int R(x) dx$?

Rewrite $R(x)$ as sum of simpler rational functions. Then use substitution rule.

Example 1. $\frac{x}{x+4} = \frac{x+4-4}{x+4} = 1 - \frac{4}{x+4}$

improper *proper*

$$\int \frac{x}{x+4} dx = \int 1 - \frac{4}{x+4} dx$$

$$= x - 4 \ln|x+4| + C$$

Example 2. $\frac{1}{x^2-4} = \frac{1}{(x-2)(x+2)}$

To decompose write

$$\frac{A}{x-2} + \frac{B}{x+2} = \frac{A(x+2) + B(x-2)}{(x-2)(x+2)}$$

Numerator gives $(A+B)x + 2(A-B) = 1$

$$\Rightarrow A = -B = \frac{1}{4}$$

$$\int \frac{1}{x^2-4} dx = \frac{1}{4} \int \frac{1}{x-2} - \frac{1}{x+2} dx$$

$$= \frac{1}{4} (\ln|x-2| - \ln|x+2|) + C$$

$$= \frac{1}{4} \ln \left| \frac{x-2}{x+2} \right| + C$$

If Q is a product of distinct linear factors

$$Q = (a_1x + b_1) \cdots (a_nx + b_n)$$

take

$$R = \frac{A_1}{a_1x + b_1} + \cdots + \frac{A_n}{a_nx + b_n}$$

Example 3.
$$\frac{5x^2+2}{x(x^2+2x+2)} = \frac{A}{x} + \frac{Bx+C}{x^2+2x+2}$$

$$\Rightarrow Ax^2 + 2Ax + 2A + Bx^2 + Cx = 5x^2 + 2$$

$$\Rightarrow \begin{cases} A+B=5 \\ 2A+C=0 \\ 2A=2 \end{cases} \Rightarrow \begin{cases} A=1 \\ B=4 \\ C=-2 \end{cases}$$

$$\int \frac{5x^2+2}{x(x^2+2x+2)} dx = \int \frac{1}{x} + \frac{4x-2}{x^2+2x+2} dx$$

$$= \ln|x| + 2 \int \frac{2x-1}{(x+1)^2+1} dx$$

$$= \ln|x| + 2 \int \frac{2(u-1)-1}{u^2+1} du \quad u=x+1$$

$$= \ln|x| + 4 \int \frac{u}{u^2-1} du - 6 \int \frac{1}{u^2+1} du$$

$$= \ln|x| + 2 \int \frac{d(u^2)}{u^2-1} - 6 \arctan u + C$$

$$= \ln|x| + 2 \ln|u-1| - 6 \arctan u + C$$

$$= 3 \ln|x| - 6 \arctan(x+1) + C$$

If Q contains distinct irreducible quadratic factors, take $\frac{Ax+B}{ax^2+bx+c}$ for the quadratic term.

Example 4
$$\frac{4x}{x^3 - x^2 - x + 1} = \frac{4x}{(x-1)^2(x+1)}$$

$$= \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1}$$

$$\Rightarrow 4x = A(x+1)(x-1) + B(x-1) + C(x-1)^2$$

$$= (A+C)x^2 + (B-2C)x + (-A+B+C)$$

$$\Rightarrow A=1 \quad B=2 \quad C=-1$$

$$\int \frac{4x}{x^3 - x^2 - x + 1} dx = \int \frac{1}{x-1} + \frac{2}{(x-1)^2} - \frac{1}{x+1} dx$$

$$= \ln|x-1| - \frac{2}{x-1} - \ln|x+1| + C$$

$$= \ln \left| \frac{x-1}{x+1} \right| - \frac{2}{x-1} + C$$

If Q contains repeated linear factor say $(ax+b)^r$
take $\frac{A_1}{ax+b} + \frac{A_2}{(ax+b)^2} + \dots + \frac{A_r}{(ax+b)^r}$

Similarly, for repeated quadratic factor $(ax^2+bx+c)^r$
take $\frac{A_1x+B_1}{ax^2+bx+c} + \dots + \frac{A_r x + B_r}{(ax^2+bx+c)^r}$

Last time $\int \frac{P(x)}{Q(x)} dx$


This time: Approximate integrals.

Motivation

In general, it is difficult to compute the antiderivative of a function and apply FTC. We then seek for an approximate value of the integral.

Recall in Calculus I, the integral is defined as limits of Riemann sums.

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \cdot \Delta x.$$



If instead of taking $n \rightarrow \infty$, we sum over a finite number of intervals,

$$\int_a^b f(x) dx \approx \sum_{i=1}^n f(x_i^*) \cdot \Delta x$$

For the finite sum above

- if $x_i^* = x_{i-1}$ left endpoint approx.
- if $x_i^* = x_i$ right endpoint approx.
- if $x_i^* = \frac{x_{i-1} + x_i}{2}$ midpoint approx.

We usually use midpoint approx. Let's write the above formula explicitly

Midpoint rule

$$\int_a^b f(x) dx \approx M_n = \Delta x (f(\bar{x}_1) + f(\bar{x}_2) + \dots + f(\bar{x}_n))$$

where $\bar{x}_i = \text{midpoints}$ $\Delta x = \frac{b-a}{n}$

Another way to approximate the integral is the

Trapezoidal rule

$$\int_a^b f(x) dx \approx T_n$$

$$= \frac{\Delta x}{2} (f(x_0) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(x_n))$$

where $\Delta x = \frac{b-a}{n}$

Note that $T_n =$

$$\left(\underbrace{\frac{f(x_0) + f(x_1)}{2}} + \frac{f(x_1) + f(x_2)}{2} + \dots + \frac{f(x_{n-1}) + f(x_n)}{2} \right) \underbrace{\Delta x}$$

area of trapezoid

Error of approx.

$$E_M = \int_a^b f(x) dx - M_n \quad E_T = \int_a^b f(x) dx - T_n$$

Error bounds: suppose $|f''(x)| \leq K$ for $a \leq x \leq b$ then

$$|E_M| \leq \frac{K(b-a)^3}{24n^2} \quad |E_T| \leq \frac{K(b-a)^3}{12n^2}$$

Similar to trapezoidal rule another rule to approximate integrals is

Simpson's rule

$$\begin{aligned} \int_a^b f(x) dx &\approx S_n \\ &= \frac{\Delta x}{3} \left(f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \dots \right. \\ &\quad \left. + 4f(x_{n-2}) + 2f(x_{n-1}) + f(x_n) \right) \end{aligned}$$

where $\Delta x = \frac{b-a}{n}$

Error $E_S = \int_a^b f(x) dx - S_n$

Error bound: suppose $|f^{(4)}(x)| \leq K$ for $a \leq x \leq b$ then

$$|E_S| \leq \frac{K(b-a)^5}{180n^4}$$

How to use these approx. in practice?

A naive example

Take $f(x) = x^2$, $1 \leq x \leq 4$ and consider to find $n \geq 1$
s.t. $|E_n| < 0.1$

$$\text{Given } |E_n| \leq \frac{K(b-a)^3}{24n^2}$$

here $K=2$ as $f''(x) \equiv 2$

$$b-a = 4-1 = 3$$

$$\Rightarrow \frac{2 \cdot 3^3}{24n^2} = \frac{54}{24n^2} \leq 0.1$$

$$\Rightarrow n \geq \sqrt{\frac{54}{24}} \approx 4.7$$

So the smallest n to take is 5.

Improper integrals.

So far we dealt with $\int_a^b f(x) dx$ for

- $x \in [a, b]$ a finite interval
- f piecewise continuous, finite

These are called proper integrals (note that this has nothing to do with proper $\frac{P(x)}{Q(x)}$).

In this section we are going to study improper integrals.

Type I

def
↓

$$\int_a^{\infty} f(x) dx := \lim_{t \rightarrow \infty} \int_a^t f(x) dx$$

$$\int_{-\infty}^b f(x) dx := \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$$

$$\begin{aligned} \int_{-\infty}^{\infty} f(x) dx &:= \int_{-\infty}^a f(x) dx + \int_a^{\infty} f(x) dx \\ &= \lim_{t \rightarrow -\infty} \int_t^a f(x) dx + \lim_{t \rightarrow \infty} \int_a^t f(x) dx \end{aligned}$$


An improper integral is convergent if the above limit exists, otherwise it is divergent.

Type II if $f \rightarrow \infty$ at some point $c \in [a, b]$

$$\int_a^c f(x) dx := \lim_{t \rightarrow c^-} \int_a^t f(x) dx$$

$$\int_c^b f(x) dx := \lim_{t \rightarrow c^+} \int_t^b f(x) dx$$

$$\int_a^b f(x) dx := \int_a^c + \int_c^b$$



Example 1. $\int_0^{\infty} e^{-x} dx$

$$= \lim_{t \rightarrow \infty} \int_0^t e^{-x} dx = \lim_{t \rightarrow \infty} [-e^{-x}]_0^t$$

$$= \lim_{t \rightarrow \infty} -e^{-t} - 1 = -1$$

Example 2 $\int_1^{\infty} \frac{1}{x^p} dx$ $p > 1$

$$= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx$$

$$= \lim_{t \rightarrow \infty} \left[\frac{x^{-p+1}}{-p+1} \right]_1^t$$

$$= \lim_{t \rightarrow \infty} \frac{1}{p-1} \left(\frac{1}{t^{p-1}} - 1 \right)$$

$$= \frac{1}{p-1}$$

Note that
• this diverges
if $p < 1$

• We can replace
1 with any real
number a .

Example 3. $\int_0^1 \frac{1}{x-1} dx$

$$= \lim_{t \rightarrow 1^-} \int_0^t \frac{1}{x-1} dx$$

$$= \lim_{t \rightarrow 1^-} (\ln|t-1| - \ln|-1|)$$

$$= \lim_{t \rightarrow 1^-} \ln|t-1|$$


$$= -\infty \quad \text{diverges.}$$

Comparison test for improper integrals.

Suppose $f(x)$, $g(x)$ are continuous, nonnegative and $f(x) \geq g(x)$ for $x \geq a$.

$$\int_a^{\infty} f(x) dx \text{ conv.} \Rightarrow \int_a^{\infty} g(x) dx \text{ conv.}$$

$$\int_a^{\infty} g(x) dx \text{ div.} \Rightarrow \int_a^{\infty} f(x) dx \text{ div.}$$



To remember

$$\int_a^{\infty} \frac{1}{x^p} dx \quad \begin{array}{l} \text{conv. } p > 1 \\ \text{div } p \leq 1 \end{array}$$

Example 4. Show the integral I is divergent

$$I = \int_1^{\infty} \frac{1+e^{-x}}{x} dx$$

Since $\frac{1+e^{-x}}{x} > \frac{1}{x}$ (as $e^{-x} > 0$) and

$$\int_1^{\infty} \frac{1}{x} dx$$

$$= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx = \lim_{t \rightarrow \infty} [\ln|x|]_1^t$$

$$= \lim_{t \rightarrow \infty} \ln t = \infty \text{ diverges.}$$

By comparison test I diverges.

More examples on comparison test

$$1. I = \int_1^{\infty} \frac{1}{\sqrt{x^6+1}} dx \text{ conv.}$$

Note that for $1 \leq x < \infty$
 $0 \leq x^6 \leq x^6+1$

$$\Rightarrow 0 \leq \sqrt{x^6} \leq \sqrt{x^6+1}$$

$$\Rightarrow \underbrace{\frac{1}{\sqrt{x^6+1}}}_g \leq \frac{1}{\sqrt{x^6}} = \underbrace{\frac{1}{x^3}}_f$$

Hence $\int_1^{\infty} \frac{1}{x^3} dx \text{ conv.} \Rightarrow I \text{ conv.}$

$$2. I = \int_2^{\infty} \frac{\cos^2(x)}{x^2} dx \text{ conv.}$$

Note that $0 \leq \cos^2(x) \leq 1$ for all x

$$\Rightarrow 0 \leq \underbrace{\frac{\cos^2(x)}{x^2}}_g \leq \underbrace{\frac{1}{x^2}}_f$$

Hence $\int_2^{\infty} \frac{1}{x^2} dx \text{ conv.} \Rightarrow I \text{ conv.}$

$$3. I = \int_3^{\infty} \frac{1}{x - e^{-x}} dx \text{ div.}$$

Since $0 < e^{-x} < x$ for $x > 3$

$$\Rightarrow 0 < x - e^{-x} < x < \infty$$

$$\Rightarrow 0 < \underbrace{\frac{1}{x}}_g \leq \underbrace{\frac{1}{x - e^{-x}}}_f$$

Hence $\int_3^{\infty} \frac{1}{x} dx \text{ div} \Rightarrow I \text{ div}$